

TRIANGULAR MAPS WITH THE CHAIN RECURRENT POINTS PERIODIC

JIŘÍ KUPKA

ABSTRACT. Forti and Paganoni [Grazer Math. Ber. **339** (1999), 125–140] found a triangular map $F(x, y) = (f(x), g_x(y))$ from $I \times I$ into itself for which closed set of periodic points is a proper subset of the set of chain recurrent points. We asked whether there is a characterization of triangular maps for which every chain recurrent point is periodic. We answer this question in positive by showing that, for a triangular map with closed set of periodic points and any positive real ε , every ε -chain from a chain recurrent point to itself may be represented as a finite union of ε -chains whose all points either are periodic or form a nontrivial ε -chain of some one-dimensional map g_x .

1. INTRODUCTION AND THE THEOREM

Denote by I the closed interval $[0, 1] \subseteq \mathbb{R}$ with the induced topology, by I^2 the Cartesian product $I \times I \subseteq \mathbb{R}^2$, by X an arbitrary compact metric space. If $A, B \subseteq X$ then \overline{A} , $\text{int}(A)$ and $\text{dist}(A, B)$ is the closure, the interior of A and the distance of A and B , respectively.

The set of continuous mappings of a compact metric space X into itself is denoted by $C(X, X)$. For $\varphi \in C(X, X)$ and $x \in X$, define inductively the n th iteration of φ by $\varphi^0(x) = x$, and $\varphi^n(x) = \varphi(\varphi^{n-1}(x))$; n is a member of positive integer set \mathbb{N} . The **orbit** $\text{Orb}(A)$ of a set $A \subseteq X$ is the set of its images under φ . The **trajectory** of x is the sequence $\{\varphi^n(x)\}_{n=0}^{\infty}$, and the set $\omega_\varphi(x)$ of limit points of the trajectory of x is the **ω -limit set** of x . Let $\omega(\varphi) = \bigcup_{x \in I} \omega_\varphi(x)$. The **α -limit point** of x is the limit point of some sequence $\{x_n\}_{n=0}^{\infty}$ such that $x_0 = x$

Received June 20, 2003.

2000 *Mathematics Subject Classification.* Primary 37E99, 37B20, 26A18, 54H20.

Key words and phrases. Triangular maps, periodic points, chain recurrent points.

Dedication to grants: GAČR 201/03/1153, MSM 192400002.

and $\varphi(x_n) = x_{n-1}$. Let $\alpha_\varphi(x)$ be the set of all α -limit points of x and let $\text{Fix}(\varphi) = \{x \in I \mid \varphi(x) = x\}$ be the set of fixed points of φ . A point x is a **periodic** point of φ , if $\varphi^p(x) = x$ for some $p \in \mathbb{N}$. The orbit of a periodic point x is a **cycle** and its cardinality is its **period**. Denote by $P(\varphi)$ the set of periodic points of φ . A point x is **nonwandering** if for every open neighborhood U of x there is an $n \in \mathbb{N}$ such that $\varphi^n(U) \cap U \neq \emptyset$. Let $\Omega(\varphi)$ denote the set of nonwandering points of φ .

Let $\varepsilon > 0$ be given and let x, y be points of X . An ε -**chain** from x to y is a finite sequence $\{x_0, x_1, \dots, x_n\}$ of points of X with $x = x_0$, $y = x_n$ and $d_X(\varphi(x_{k-1}), x_k) < \varepsilon$ for $k = 1, 2, \dots, n$. An ε -chain E is **nontrivial** if at least one point of E is nonperiodic. A point x is **chain recurrent** if and only if, for every $\varepsilon > 0$, there is an ε -chain from x to itself ($x \rightarrow_\varepsilon x$). Let $CR(\varphi)$ denote the set of all chain recurrent points of φ .

The relations between the above-mentioned sets are given by the next proposition which can be found, e.g., in [1].

Theorem 1.1. *If X is a compact metric space and $\varphi \in C(X, X)$ then*

$$(1) \quad \text{Fix}(\varphi) \subseteq P(\varphi) \subseteq \omega(\varphi) \subseteq \Omega(\varphi) \subseteq CR(\varphi).$$

The mentioned sets are equal if $f \in C(I, I)$ satisfies some assumptions; a long list of such assumptions can be found, e.g. in [4]. The next proposition which can be found, e.g. in [3], displays a few of them.

Theorem 1.2. [3] *$f \in C(I, I)$. Then the following conditions are equivalent:*

- (i) $P(f) = \overline{P(f)}$.
- (ii) $P(f) = \Omega(f)$.
- (iii) $P(f) = CR(f)$.

Our paper is devoted to **triangular maps of the square**, i.e., continuous maps $F : I^2 \rightarrow I^2$ of the form $F(x, y) = (f(x), g_x(y))$; the function f is called the **base** function of F . Denote by $T(I^2, I^2)$ the set of triangular maps $I^2 \rightarrow I^2$ and by d the max metric on I^2 . In 1990 L. S. Efremova proved the next result (in a more strong form).

Theorem 1.3. [2] *Let $F \in T(I^2, I^2)$ and $P(F)$ be closed. Then $P(F) = \Omega(F)$.*

Let (i)–(iii) be the properties displayed in Proposition 1.2 considered for a triangular map F . Since the set $\Omega(F)$ is closed, the Proposition 1.3 and (1) give (i) \Leftrightarrow (ii). Moreover, Forti and Paganoni in [3] found a triangular map F with closed set of the periodic points such that $CR(F) \setminus \omega(F) \neq \emptyset$. By (1), $\omega(F) \subseteq \Omega(F) \subseteq CR(F)$, so there is a problem to characterize triangular maps F with closed set $P(F)$ such that $P(F) = CR(F)$. The wanted characterization follows from the next theorem which is the main result of our paper.

Definition 1.4. Let $z_1 = (x, y_1), z_2 = (x, y_2)$ be periodic points of F . The point z_1 is **accessible** from z_2 ($z_2 \rightarrow^a z_1$) if there is an ε -chain from z_2 to z_1 by the map F restricted to $I_{Orb(x)} = \bigcup_{y \in Orb(x)} \{y\} \times I$ for any $\varepsilon > 0$. The point z_1 is **nontrivially accessible** from z_2 ($z_2 \rightarrow^n z_1$) if $z_2 \rightarrow^a z_1$ and for any sufficiently small $\varepsilon > 0$ any ε -chain from z_2 to z_1 by the map F restricted to $I_{Orb(x)}$ is nontrivial.

Let K_1, K_2 be subsets of $P(F)$. Then K_1 is **accessible** from K_2 ($K_2 \rightarrow^a K_1$) if $z_2 \rightarrow^a z_1$ for some $z_1 \in K_1$ and $z_2 \in K_2$.

Definition 1.5. Points $z_1, z_2 \in P(F)$ form a **t-pair** if (i) $z_2 \rightarrow^n z_1$ and, for any $\delta > 0$, (ii) there exists a finite number of connected components $K_i, i = 1, 2, \dots, m$, of $P(F)$ such that $z_1 \in K_1, z_2 \in K_m$ and, for all $i = 1, 2, \dots, m - 1$, either $K_i \rightarrow^a K_{i+1}$ or $\text{dist}(K_i, K_{i+1}) < \delta$.

Theorem 1.6. Main theorem *Let $F \in T(I^2, I^2)$ and $P(F)$ be closed. Then $CR(F) \setminus P(F) \neq \emptyset$ if and only if there exists a t-pair.*

2. KNOWN FACTS

For a map $F \in T(I^2, I^2)$, let $A(F)$ Denote any of the sets $\text{Fix}(F), P(F), \omega(F), \Omega(F)$ and $CR(F)$ and let π be the canonical projection $(x, y) \mapsto x$.

Theorem 2.1. [4] *Let $F \in T(I^2, I^2)$, let f be the base function of F . Then $\pi(A(F)) = A(f)$.*

We already know that $CR(\varphi)$ is closed for any $\varphi \in C(X, X)$. Of course, $CR(\varphi)$ enjoy a number of other general properties used later.

Theorem 2.2. [1] *Let $\varphi \in C(X, X)$. Then, for any $n \in \mathbb{N}$, $CR(\varphi) = CR(\varphi^n)$ and $P(\varphi) = P(\varphi^n)$.*

Theorem 2.3. [1] *Let $\varphi \in C(X, X)$. Then $CR(\varphi) = CR(\varphi|_{CR(\varphi)})$, i.e. every chain recurrent point remains chain recurrent for the restriction of φ to $CR(\varphi)$.*

Let $\varphi \in C(X, X)$. A nonempty closed set A contained in X is **Lyapunov stable** if, for each open set U containing A , there exists an open set V containing A such that $\varphi^n(V) \subseteq U$ for all $n \in \mathbb{N}$. A nonempty closed set A is an **attractor** if there exists an open set B containing A such that $\omega_\varphi(x) \subseteq A$ for every $x \in B$. If a nonempty closed set A is both Lyapunov stable and an attractor, we say that A is **asymptotically stable**.

Theorem 2.4. [1] *Let $\varphi \in C(X, X)$. If $A \subseteq X$ is an asymptotically stable set, then there exists an open set W containing A such that $\varphi(\overline{W}) \subseteq W$. Moreover, for any open upset U of A we can choose W so that $\overline{W} \subseteq U$.*

Denote by $Q(x, \varphi)$ the intersection of all asymptotically stable sets containing $\omega_\varphi(x)$.

Theorem 2.5. [1] *Let $\varphi \in C(X, X)$. If $y \in Q(x, \varphi)$, then $Q(y, \varphi) \subseteq Q(x, \varphi)$.*

Theorem 2.6. [1] *Let $\varphi \in C(X, X)$. If $y \in Q(x, \varphi)$, then there is an ε -chain from x to y by φ for any $\varepsilon > 0$.*

3. LEMMAS AND PROOFS

Let $\varphi \in C(X, X)$ and $K, L \subseteq X$. Denote by $S_\varepsilon(K, L)$ the closure of $\{z \in X \mid z \text{ is a member of an } \varepsilon\text{-chain from a point of } K \text{ to a point of } L \text{ by } \varphi\}$. If $K \xrightarrow{\varepsilon} L$ for any $\varepsilon > 0$, the set $S_\varphi(K, L) = \bigcap_{\varepsilon > 0} S_\varepsilon(K, L)$ is clearly nonempty and closed. If $z_0 \in CR(\varphi)$, the set $S_\varphi(z_0, z_0)$ is the **independent set** of z_0 .

Lemma 3.1. *Let $\varphi \in C(X, X)$ and $z_0 \in CR(\varphi)$. Then the independent set $S := S_\varphi(z_0, z_0)$ is strongly invariant, i.e. $\varphi(S) = S$.*

Proof. This easily follows from the continuity of φ and the definition of S . □

Lemma 3.2. *Let $\varphi \in C(X, X)$ and $z_0 \in CR(\varphi)$. Then any point of the independent set $S := S_\varphi(z_0, z_0)$ is chain recurrent for the restriction of φ to S , i.e. $S = CR(\varphi|_S)$.*

Proof. It is easy to see, $CR(\varphi|_S) \subseteq S$. We prove the converse inclusion.

Clearly $S \subseteq CR(\varphi)$. Let z be a chain recurrent point lying in S . At first we show that for any open neighborhood U of S and any $\varepsilon > 0$ there is an ε -chain from z to z in U . Assume, contrary to what we wish to show, that there exists an open neighborhood U of S and $\varepsilon > 0$ such that every ε -chain from z to itself contains a point in the complement of U . Let $\{\varepsilon_n\}$ be a decreasing sequence of positive numbers tending to 0. Then, for each $n \in \mathbb{N}$, there is a point z_n of an ε_n -chain lying outside U . Denote by \tilde{z} an accumulation point of $\{z_n\}_{n \in \mathbb{N}}$. Evidently, $\tilde{z} \notin U$. But $z \rightarrow_\varepsilon \tilde{z}$ and $\tilde{z} \rightarrow_\varepsilon z$ for every $\varepsilon > 0$ and, by the definition of S and $z \in S$, this implies $\tilde{z} \in S$, which is a contradiction.

It remains to show that for any $\varepsilon > 0$ there is an ε -chain from z to itself within S . By the continuity of φ , we can choose $\delta > 0$, $0 < \delta < \frac{\varepsilon}{3}$, so that $d_X(x, y) < \delta$ implies $d_X(\varphi(x), \varphi(y)) < \frac{\varepsilon}{3}$. Let U be an open δ -neighborhood of S and let $\{z_0, z_1, \dots, z_k\}$ be an $\frac{\varepsilon}{3}$ -chain from z to z lying in U . There exist points s_0, s_1, \dots, s_k in S with $s_0 = s_k = z$ and $d_X(z_j, s_j) < \delta$ for $1 \leq j \leq k$. Then for each $j = 1, 2, \dots, k$

$$d_X(\varphi(s_{j-1}), s_j) < d_X(\varphi(s_{j-1}), \varphi(z_{j-1})) + d_X(\varphi(z_{j-1}), z_j) + d_X(z_j, s_j) < \varepsilon,$$

which already proves $S = CR(\varphi|_S)$. □

Lemma 3.3. *Let $F \in T(I^2, I^2)$, $P(F)$ be closed and $z = (x, y) \in CR(F) \setminus P(F)$. If x is a fixed point of the base map f , then the independent set $S_f(x, x)$ is a nondegenerate closed interval of $P(f)$.*

Proof. Let the assumptions be fulfilled. Clearly,

$$(2) \quad \pi(S_F(z, z)) \subseteq S_f(x, x).$$

Denote $S := S_F(z, z)$ and $D := S_f(x, x)$. Since any independent set is closed, strongly invariant and the fixed point x belongs to D , D is closed interval.

We show that D is nondegenerate. Assume, on the contrary, $D = \{x\}$. By (2), $S \subseteq \{x\} \times I$ and hence, with respect to Lemma 3.2, $y \in CR(g_x)$ since $z \in S$. On the other hand, $z \notin P(F)$ implies $y \notin P(g_x)$, i.e. $y \in CR(g_x) \setminus P(g_x)$. So, by Proposition 1.2, $P(g_x)$ is not closed, which contradicts $P(F) = \overline{P(F)}$.

Finally, we show $D \subseteq P(f)$. By Lemma 3.2, any point of D is chain recurrent. Recall that $P(F)$ is closed and the projection of any closed set is closed. Hence, by Propositions 1.2 and 2.1, $D \subseteq P(f)$. \square

Lemma 3.4. *Let $F \in T(I^2, I^2)$, $P(F)$ be closed and f be the base map of F . Let $z = (x, y) \notin P(F)$ be such that $x \in \text{Fix}(f)$. If any $z_1 \in \omega_F(z)$, $z_2 \in \alpha_F(z)$ lie in the same connected component of $P(F)$, then z_1, z_2 form a t-pair. Otherwise $z_2 \xrightarrow{n} z_1$.*

Proof. Let the assumptions be fulfilled. Clearly, $z_1 = (x, y_1), z_2 = (x, y_2)$ are periodic points of F . We show that $y_1 \neq y_2$. Really, by the choice of z_1, z_2 , for any $\varepsilon > 0$, $y \rightarrow_\varepsilon y_1$ and $y_2 \rightarrow_\varepsilon y$ by g_x . Hence $y_1 = y_2$ implies $y \in CR(g_x)$ and, accordingly, $CR(g_x) \setminus P(g_x) \neq \emptyset$. So, by Proposition 1.2, $P(g_x)$ is not closed, a contradiction.

It is easy to see that $z_2 \xrightarrow{a} z_1$. Hence z_1, z_2 form a t-pair if they lie in the same connected component of $P(F)$. In the opposite case, $z_2 \xrightarrow{n} z_1$. \square

Proof of the main theorem. Let F be a triangular map and f its base map. The existence of a t-pair clearly implies the existence of nonperiodic chain recurrent point. Let us prove the converse implication.

Let $z_0 = (x_0, y_0)$ be a nonperiodic chain recurrent point, $S := S_F(z_0, z_0)$ and $D := S_f(x_0, x_0)$. We show that there is a t-pair in S . Clearly, any t-pair of some iteration of F is t-pair of F . Moreover, by Proposition 2.2, $z_0 \in CR(F^n) \setminus P(F^n)$ for any $n \in \mathbb{N}$. So we may consider some convenient iteration of F instead of F . Thus we may assume $x_0 \in \text{Fix}(f)$, since, by Propositions 1.2 and 2.1, $x_0 \in P(f)$. Obviously, $S \subseteq D \times I$, where, by Lemma 3.3, D is a closed nondegenerate interval of $P(f)$. But any interval of $P(f)$ containing a fixed point has only periodic points of period ≤ 2 . So we find more convenient iteration of F and may assume $\pi(S) \subseteq \text{Fix}(f)$. Moreover, by Lemma 3.2, $z_0 \in CR(F|_S) \setminus P(F|_S)$.

In the rest of the proof put $F := F|_S$ and $P := P(F) \cap S$. Since the complement of P is open, there is a $\delta > 0$ such that an open δ -neighborhood of z_0 contains no periodic point. Since S is strongly invariant, P is nonempty and hence we can define a relation $\sim := \sim(\delta)$ on the set of all connected components of P in the following way: $K \sim L$ if and only if $K \rightarrow_\delta L$ by F restricted to P . It is easy to see that \sim is an equivalence relation on P . Members of the decomposition $P|_\sim$ are **δ -components**. The distance $d(v, w) \geq \delta$ if $v \in K, w \in L$ are points of distinct δ -components K, L and hence, by the compactness of S , there is only a finite number of δ -components $H_i, i = 1, 2, \dots, k$.

Lemma 3.5. *Let H be a δ -component and $u, w \in H$. Then there are connected components L_1, L_2, \dots, L_r of H such that $u \in L_1, w \in L_r$ and $\text{dist}(L_i, L_{i+1}) < \delta$ for every $i = 1, 2, \dots, r - 1$.*

Proof. Obvious. □

By Lemma 3.1, $\omega_F(z_0) \subseteq P$ and $\alpha_F(z_0) \cap P \neq \emptyset$ since $z_0 \in S$. Choose $z_1 \in \omega_F(z_0)$. We show that there is a $z_2 := z_2(\delta) \in \alpha_F(z_0) \cap P$ such that z_1 and z_2 satisfy the condition (ii) of Definition 1.5 ($z_2 \xrightarrow{\delta} z_1$). With respect to Lemma 3.4, we may assume $z_2 \xrightarrow{n} z_1$. By Lemma 3.5, it is sufficient to find δ -components K_1, K_2, \dots, K_m such that $z_1 \in K_1, z_2 \in K_m$ and

$$(3) \quad K_i \xrightarrow{a} K_{i+1} \text{ for } i = 1, 2, \dots, m - 1.$$

Without loss of generality, assume $z_1 \in H_1$. If $\alpha_F(z_0) \cap H_1 \neq \emptyset$, by Lemmas 3.4 and 3.5, $z_2 \xrightarrow{\delta} z_1$ for every z_2 from this intersection. So let $\alpha_F(z_0) \cap (P \setminus H_1) \neq \emptyset$ and $K_1 := H_1$. Clearly $K \xrightarrow{a} L$ implies $\pi(K) \cap \pi(L) \neq \emptyset$. Hence we may restrict our attention to the set $S_1 := S \cap (\pi(K_1) \times I)$ to show that there is a δ -component H such that $K_1 \xrightarrow{a} H$.

Lemma 3.6. $K_1 \xrightarrow{a} H$ for some $H \in P|_\sim$.

Proof. Assume, on the contrary, that

$$(4) \quad K_1 \not\xrightarrow{a} H \text{ for each } H \in P|_\sim \setminus K_1.$$

Fix $x \in \pi(S_1) \subseteq \text{Fix}(f)$. By Proposition 2.6, $Q(y, g_x) \cap (P \setminus K_1) = \emptyset$ for any $(x, y) \in K_1(x) := K_1 \cap I_x$. Hence $Q(x) \cap P = K_1$, where $Q(x)$ denotes $\bigcup_{(x,y) \in K_1(x)} Q(y, g_x)$. It is well known fact, that $Q(\cdot, \cdot)$ is an asymptotically stable set. Generally, only a finite union of asymptotically stable sets must be asymptotically stable, see [1]. Consequently, it suffices to clarify that $Q(x)$ is "generated" by a finite number of points of $K_1(x)$ to show that $Q(x)$ is asymptotically stable.

Since $P(F)$ is closed, K_1 is closed and hence, any sequence $\{y_i\} \subseteq K_1$ has an limit point $y \in K_1(x)$. Because $Q(y, g_x)$ is asymptotically stable, there exists $N \in \mathbb{N}$ such that $y_n \in Q(y, g_x)$ for every $n \geq N$. Moreover, if J is subinterval of $K_1(x)$, $J \subseteq Q(y, g_x)$ for any $(x, y) \in J$. By these two facts, Proposition 2.5 and the compactness of $K_1(x)$, $Q(x)$ is asymptotically stable.

Accordingly, by (4), $Q(x)$ is asymptotically stable for any $x \in \pi(S_1)$. Since we have restricted F to S_1 , there is, by Proposition 2.4, an open set W containing K_1 such that

$$(5) \quad F(\overline{W}) \subseteq W.$$

Clearly, again by Proposition 2.4, we may assume $z_0 \notin W$. But we have shown that $\omega_F(z_0) \subseteq K_1$. Hence, for a sufficiently small $\varepsilon > 0$, some point of any ε -chain from z_0 to itself lies in W . On the other hand, since W is a subset of any open neighborhood of K_1 , by (5), there is no ε -chain from H_1 to z_0 for some sufficiently small $\varepsilon > 0$, a contradiction. \square

Denote $H'_1 := K_1$. By Lemma 3.6, a set $H'_2 = \{H \in P|_{\sim} \setminus H'_1 \mid K_1 \rightarrow^a H\}$ is nonempty. If $\alpha_F(z_0) \cap H'_2 \neq \emptyset$, by Lemmas 3.4 and 3.5, there is a $z_2 := z_2(\delta) \in \alpha_F(z_0) \cap H'_2$ such that $z_2 \rightarrow_{\delta}^t z_1$. If $\alpha_F(z_0) \cap H'_2$ is empty, we analogously define the set $H'_3 = \{H \in P|_{\sim} \setminus (H'_1 \cup H'_2) \mid \exists L \in H'_2 : L \rightarrow^a H\}$ and similarly show that $H'_3 \neq \emptyset$. Accordingly, since the cardinality of $P|_{\sim}$ is finite, there are nonempty sets H'_1, H'_2, \dots, H'_m , $m \leq k$, of δ -components of P such that $\alpha_F(z_0) \cap K \neq \emptyset$ for some δ -component $K \in H'_m$ and, for any $j \in \{1, 2, \dots, m-1\}$, any member of H'_{j+1} is accessible from some member of H'_j .

This already proves that for any $\delta > 0$ there is a $z_2 := z_2(\delta) \in \alpha_F(z_0) \cap S$ such that $z_2 \xrightarrow{t_\delta} z_1$. It is easy to see that $z_2(\delta) \xrightarrow{t_\tau} z_1$ for any $\tau > \delta$. Accordingly, for any decreasing sequence of positive reals $\{\delta_i\}$ tending to 0 the limit point \tilde{z}_2 of the sequence $\{z_2(\delta_i)\}$ and z_1 is a t-pair. \square

1. Block L. S. and Coppel W. A., *Dynamics in One Dimension*, Lecture Notes in Math. 1513, Springer, Berlin – Heidelberg – New York, 1992.
2. Efremova L. S., *On the nonwandering set and the center of triangular maps with closed set of periodic points in the base* (in Russian), *Dynam. Systems and Nonlinear Phenomena*, Inst. of Mathem. of NAS of Ukraine, Kiev (1990), 15–25.
3. Forti G. L. and Paganoni L., *On some properties of triangular maps*, *Grazer Math. Ber.* **339** (1999), 125–140.
4. Kolyada S. F., *On dynamics of triangular maps of the square*, *Ergodic Theory Dynam. Systems* **12** (1992), 749–768.

Jiří Kupka, Matematický ústav v Opavě, Slezská univerzita v Opavě, Na Rybníčku 1, 746 01 Opava, Czech Republic, *e-mail*: Jiri.Kupka@math.slu.cz,
<http://www.math.slu.cz/People/JiriKupka.php>