

ON THE FRAGMENTAL STRUCTURES

M. A. KAMAL AND N. S. MAHMOUD

ABSTRACT. In this work we study the fragment structures over a ring extension R of a ring R_0 . The defining conditions of the fragments with the partial actions on the descending chains of R_0 -modules measure how far they are from being R -modules. The category of R -fragments lies between the categories of R_0 -modules and of R -modules. In spite of R -fragments, in a general setting, are far from being R -modules; they behave, in some ways, the same as R -modules. We prove some important results for finitely spanned fragments and some of their related properties.

1. INTRODUCTION

Let R be a ring with unity 1. Let $R_0 \subseteq R_1 \subseteq \dots \subseteq R$ be a positive filtration of R ; with $1 \in R_0$. An abelian group $(M, +)$ with the descending chain

$$M = M_0 \supseteq M_1 \supseteq M_2 \supseteq \dots \supseteq M_n \supseteq \dots$$

of subgroups of M is called a left R -fragment with respect to the mappings $\varphi_{i,j} : R_i \times M_j \rightarrow M_{j-i}$ (for all i, j ; with $j \geq i$) if the following conditions are satisfied:

1. $\varphi_{i,j}|_{R_q \times M_r} = \varphi_{q,r}$; for all $q \leq i \leq j \leq r$.
2. $\varphi_{i,j}(\alpha, m+n) = \varphi_{i,j}(\alpha, m) + \varphi_{i,j}(\alpha, n)$; for all $\alpha \in R_i, m, n \in M_j$, and $i \leq j$.
3. $\varphi_{i,j}(\alpha + \beta, m) = \varphi_{i,j}(\alpha, m) + \varphi_{i,j}(\beta, m)$; for all $\alpha, \beta \in R_i, m \in M_j$, and $i \leq j$.
4. $\varphi_{i,i}(\alpha, \varphi_{i,(i+j)}(\beta, m)) = \varphi_{(i+j),(i+j)}(\alpha\beta, m)$; for all $\alpha \in R_i, \beta \in R_j$, and $m \in M_{i+j}$.
5. $\varphi_{i,j}(1, m) = m$; for all $m \in M_j$.

Received February 23, 2003.

2000 *Mathematics Subject Classification.* Primary 15A87.

A right R -fragment can be defined in a similar fashion. For all $\alpha \in R_i$ and $m \in M_j$, we write αm for $\varphi_{i,j}(\alpha, m)$. We shall usually say simply that M , rather than $M = M_0 \supseteq M_1 \supseteq M_2 \supseteq \dots \supseteq M_n \supseteq \dots$ with respect to $\varphi_{i,j} : R_i \times M_j \rightarrow M_{j-i}$ (for all i, j ; with $j \geq i$), is a left R -fragment. This allows some ambiguity, for a given chain of abelian groups may admit more than one left R -fragment structure. We clear this out by fixing a chain of subgroups of M , $M_0 \supseteq M_1 \supseteq M_2 \supseteq \dots \supseteq M_n \supseteq \dots$ and certain maps $\varphi_{i,j} : R_i \times M_j \rightarrow M_{j-i}$.

Observe that, any left R -module M can be considered as a left R -fragment with respect to the chain $M = M_0 \supseteq M_1 \supseteq M_2 \supseteq \dots \supseteq M_n \supseteq \dots$, where $M_i = M$ for all i , and $R_0 \subseteq R_1 \subseteq \dots \subseteq R$ is any positive filtration of R , with rm taken to be the left multiplication of elements of R by elements of M as a left R -module; for all $r \in R_i$, $m \in M_j = M$. Observe also that, each abelian subgroup M_i , in the chain of a left R -fragment M , is a left R_0 -module.

Let M be a left R -fragment, we denote $\bigcap_{i \in I} M_i$ by $B(M)$ and call it the body of M . If the filtration of R is exhaustive; i.e. $R = \bigcup_{i \in I} R_i$, then the left actions of the elements of R_i on the elements of M_j (for all i, j) induce an R -module structure on $B(M)$. It is clear that $B(M)$ is the largest abelian subgroup of M on which the left actions of R_i on M_j form a left R -module structure.

In this paper, R will be a filtered ring with a positive exhaustive filtration $\{R_i\}_{i \in I}$; i.e. $R_0 \subseteq R_1 \subseteq \dots \subseteq R$, and $R = \bigcup_{i \in I} R_i$.

Let M be a left R -fragment. We simply say X is subset of M , for any subset X of M_0 ; and we say x is an element of M for any element x of M_0 . Let $x \in M$ be an arbitrary element, then either $x \in M_n$ and $x \notin M_{n+1}$ for some positive integer n , or $x \in B(M)$. In the first case x is called of depth n (denoted by $d_M(x) = n$), while in the second case we say x has an infinite depth (i.e. $d_M(x) = \infty$). It is clear that $d_M(0) = \infty$.

Let $N = N_0 \supseteq N_1 \supseteq N_2 \supseteq \dots \supseteq N_n \supseteq \dots$ be a chain of subgroups of a left R -fragment M . N is called a subfragment of M if $N_i \subseteq M_i$ ($i = 1, 2, \dots$), and for all i, j with $j \geq i$, $rx \in N_{j-i}$ for all $r \in R_i$ and $x \in N_j$. If $N_i = 0$ for all i , then N is called the zero subfragment, and is denoted by 0 .

2. NONTRIVIAL SUBFRAGMENTS

A subfragment N of a left R -fragment M is called a nontrivial subfragment in case of if $R_i x \subseteq N_0$, for $x \in N_0$ with $d_M(x) \geq i$, then $x \in N_i$. N is called an improper (otherwise it is called proper) subfragment of M if, for some $j \geq 0$, $N_0 = M_j$, for each $i \geq 1$ there exists $k(i) \geq 1$ such that $N_i = M_{j+k(i)}$.

Lemma 1. *Let M be a left R -fragment such that $M_i = M_{i+1}$ for all $i = 0, 1, \dots$. If N is a nontrivial subfragment such that N_0 is an R -submodule of M , then $N_i = N_{i+1}$ for all i .*

Corollary 2. *Let M be a left R -fragment such that the filtration of R is given by $R_0 = R_1 = \dots = R$, and $M_i = M_{i+1}$ for all $i = 0, 1, \dots$; i. e. M is a left R -module. Then the chain of any nontrivial subfragment N of M must be of the form $N_0 = N_1 = N_2 = \dots = N_n = \dots$ for some R -submodule N_0 of M .*

Lemma 3. *Let N be a subfragment of a left R -fragment M such that $N_0 = M_0$. If N is nontrivial, then $N_i = M_i$ for all i ; i. e. $N = M$.*

Proof. If $x \in M_k$, then $R_k x M_0 = N_0$. This yields $x \in N_k$, and hence $M_k = N_k$ for all k . □

Let M be a left R -fragment, and $\{N^i\}_{i \in I}$ be a family of subfragments of M . Then the chain

$$\bigcap_{i \in I} N_0^i \supseteq \bigcap_{i \in I} N_1^i \supseteq \dots \supseteq \bigcap_{i \in I} N_n^i \supseteq \dots$$

of subgroups of M forms a subfragment. This subfragment of M is called the fragment intersection of the family $\{N^i\}_{i \in I}$, and is denoted by $\bigcap_{i \in I} N^i$. Observe that

$$\bigcap_{i \in I} N^i := \bigcap_{i \in I} N_0^i.$$

Let $\{N^i\}_{i \in I}$ be a chain of subfragments of M , and consider the following chain

$$\bigcap_{i \in I} N_0^i \supseteq \bigcap_{i \in I} N_1^i \supseteq \dots \supseteq \bigcap_{i \in I} N_k^i \supseteq \dots$$

of subgroups of M . It is clear that it forms a chain of a subfragment of M . This subfragment is called the fragment union of the chain $\{N^i\}_{i \in I}$.

Lemma 4. *The intersection of nontrivial subfragments of a left R -fragment is nontrivial.*

Proof. Let $\{N^i\}_{i \in I}$ be a family of nontrivial subfragments of a left R -fragment M , and let $x \in \bigcap_{i \in I} N_0^i$. Then the nontriviality of each N^i yeiles $x \in \bigcap_{i \in I} N_j^i$, whenever $R_j x \subseteq \bigcap_{i \in I} N_0^i$ and $d_M(x) \geq j$. \square

3. SUBFRAGMENT SPANNED BY A SUBSET

Let M be a left R -fragment, and X be a subset of M . Define the subset $L(X)$ of M by:

$$L(X) = \{r_n(r_{n-1}(\dots(r_0 x))) : x \in X, \text{ for some } n, r_i \in R_i, \\ \text{and } i \geq d_M(r_{i-1}(\dots(r_0 x))); i = 0, 1, \dots, n\}.$$

A subfragment N of M is said to be strictly containing X if $L(X) \subseteq N$.

Lemma 5. *Let $\{X_i\}_{i \in I}$ be a family of subsets of a left R -fragment M . Then:*

- 1) $X_i \subseteq L(X_i)$, and if $X_i \subseteq X_j$, then $L(X_i) \subseteq L(X_j)$.
- 2) $L(\bigcup_{i \in I} X_i) = \bigcup_{i \in I} L(X_i)$.

Proof is clear.

Lemma 6. *Let X be a subset of a left R -fragment M , then the intersection of all subfragments, which are strictly containing X , is strictly containing X .*

Proof is clear.

Let Γ be the set of all nontrivial subfragments of M which are strictly containing a subset X . By Lemma 4, and Lemma 6, the intersection $\bigcap \Gamma$ is again a nontrivial subfragment of M , which is strictly containing X . We call it the subfragment spanned by X , and is denoted by $\prec X \succ$. If X is the empty set, then by convention $\prec X \succ = 0$.

Lemma 7. *Let M be a left R -fragment, and X be a subset of M . Define*

$$N_0 = \left\{ \sum_{i=1}^n t_i : t_i \in L(X), n \text{ is a positive integer} \right\},$$

$$N_k = \{y \in N_0 : d_M(y) \geq k, \text{ and } R_k y \subseteq N_0\}; \quad k = 1, 2, \dots .$$

Then

$$N := N_0 \supseteq N_1 \supseteq N_2 \supseteq N_n \supseteq \dots$$

is a subfragment of M , and the subfragment spanned by X is just N .

Proof. It is clear that, from the definition of N_k , that

$$N := N_0 \supseteq N_1 \supseteq N_2 \supseteq N_n \supseteq \dots$$

is a descending chain of abelian subgroups of M . Now let $y \in N_k$ and $r_i \in R_i$, where $i \leq k$. Since $d_M(y) \geq k$, we have that $r_i y \in M_{k-i}$; and hence $d_M(r_i y) \geq k - i$. We have

$$R_{k-i}(r_i y) \subseteq R_{k-1}(R_i y) \subseteq R_k y \subseteq N_0;$$

and thus $r_i y \subseteq N_{k-i}$. This shows that N is a subfragment of M . It is clear, from the definition of N_0 , that $L(X) \subseteq N_0$; i.e. N is strictly containing X . It is also clear, from the definition of N_i , that N is nontrivial. Therefore $\prec X \succ \subseteq N$. To show that $N \subseteq \prec X \succ$, let K be any nontrivial subfragment which is strictly containing X . Since $L(X) \subseteq K$, we have that N_0 is contained in K_0 . Now let $y \in N_m$, for some positive integer m . Hence $R_m y \subseteq N_0 \subseteq K_0$, and thus the nontriviality of K yields $y \in K_m$. Therefore $N \subseteq \prec X \succ$. \square

If X is a subset of a left R -fragment M such that $\prec X \succ = M$, then X is said to span M , and X is called a spanning set for M . M with a finite spanning set is said to be finitely spanned. M with a single element spanning set is a cyclic fragment.

Corollary 8. *If X is a spanning set for a left R -fragment M , then the chain of M is given by :*

$$M_k = \left\{ \sum_{i=1}^n y_i : y_i \in L(X), d_M\left(\sum_{i=1}^n y_i\right) \geq k, n \text{ is a positive integer} \right\}; \quad k = 0, 1, \dots$$

Corollary 9. *If X is a spanning set for a left R -fragment M , with $d_M(x) = \infty$ for all $x \in X$, then $M_i = M_{i+1}$ for all i ; i.e. the fragment structure forms a left R -module structure on M .*

A nontrivial subfragment K of a left R -fragment M is said to be a strict subfragment if K is strictly containing K_0 . By Lemma 5, we have that $L(X) \subseteq K_0$ for any subset X of K_0 , and that $L(K_0) = K_0$.

Lemma 10. *The following are equivalent for a nontrivial subfragment K of M :*

- 1) K is strict,
- 2) $\prec X \succ$ is contained in K , for any subset X of K ,
- 3) $K_i = M_i \cap K_0$, for all $i = 0, 1, 2, \dots$,
- 4) $d_M(x) = d_K(x)$, for all $x \in K$.

Proof. 1) \Rightarrow 2) is clear.

2) \Rightarrow 3) It is clear that, for all i , $K_i \subseteq M_i \cap K_0$. Now let $x \in M_i \cap K_0$. Then by 2), $R_i x \subseteq \prec x \succ \subseteq K_0$. But since K is nontrivial, we have that $x \in K_i$. Hence $K_i = M_i \cap K_0$.

3) \Rightarrow 4) It is clear that $d_K(x) \leq d_M(x)$ for all $x \in K$. Now let $d_M(x) = n$, $x \in K$, then $x \in K_0 \cap M_n = K_n$, and $x \notin K_{n+1}$ (due to $K_{n+1} \subseteq M_{n+1}$); i.e. $d_K(x) = n$. It is clear that $d_K(x) = \infty$ whenever $d_M(x) = \infty$.

4) \Rightarrow 1) Let X be a subset of K . By 4), it follows that $L(X) \subseteq K$; i.e. K is strictly containing X for all subsets X of K . Hence we have 1). □

Remarks. 1) From Corollary 8, it is clear that a left R -module M is finitely generated if and only if it is finitely spanned as a natural left R -fragment.

2) If M is a left R -fragment, and if the factor R_0 -module M_n/M_{n+i} is finitely generated for all $n = 0, 1, \dots$, then M need not be a finitely spanned fragment. In fact $M_n/M_{n+i} = 0$ is finitely generated for every R -module M , and for all $n = 0, 1, \dots$

3) The intersection of an arbitrary family of strict subfragments of a left R -fragment is strict.

Let M be a left R -fragment, and let $\{N^i\}_{i=1}^m$ be a family of subfragments of M , where the chain of N^i is given by

$$N_0^i \supseteq N_1^i \supseteq N_2^i \supseteq \dots \supseteq N_n^i \supseteq \dots;$$

$i = 1, 2, \dots, m$. Then the following chain

$$\sum_{i=1}^m N_0^i \supseteq \sum_{i=1}^m N_1^i \supseteq \dots \supseteq \sum_{i=1}^m N_n^i \supseteq \dots$$

forms a subfragment of M (denoted by $\sum_{i=1}^m N^i$) called the fragment sum of the family $\{N^i\}_{i=1}^m$. If each N^i is

a strict subfragment, then we define the fragment sum $\sum_{i=1}^m N^i$ to be the subfragment of M spanned by the subset

$\bigcup_{i=1}^m N_0^i$. It is clear, by Lemma 5, that

$$L\left(\bigcup_{i=1}^m N_0^i\right) = \bigcup_{i=1}^m L(N_0^i) = \bigcup_{i=1}^m N_0^i.$$

Hence $\left(\sum_{i=1}^m N^i\right)_0 = \langle \bigcup_{i=1}^m N_0^i \rangle_0 = \sum_{i=1}^m N_0^i$, whenever each of N^i is strict.

Remark. If $\{N^i\}_{i \in I}$ is an arbitrary collection of strict subfragments of a left R -fragment M , then motivated by the finite sum of strict subfragments, we define the fragment sum $\sum_{i \in I} N^i$ of $\{N^i\}_{i \in I}$ to be the subfragment of M spanned by $\bigcup_{i \in I} N_0^i$.

Lemma 11. *Let $\{N^i\}_{i \in I}$ be an arbitrary collection of strict subfragments of a left R -fragment M . Then*

$$K := \bigcup \left\{ \sum_{i \in F} N^i : F \text{ is a finite subset of } I \right\},$$

with the chain

$$K_0 \supseteq K_1 \supseteq K_2 \supseteq \dots \supseteq K_n \supseteq \dots,$$

where

$$K_n = \bigcup \left\{ \left(\sum_{i \in F} N^i \right)_n : F \text{ is a finite subset of } I \right\},$$

is a strict subfragment of M .

Proof. It is clear that K with the given chain is a subfragment of M . Let $x \in K_0$ such that $R_j x \subseteq K_0$, and $j \leq d_M(x)$. Then $x \in \left(\sum_{i \in F} N^i \right)_0$ for some finite subset F of I . Since $\sum_{i \in F} N^i$ is a strict subfragment of M , we have by Lemma 10 that $d_{N(F)}(x) = d_M(x) = m \geq j$, where $N(F) := \sum_{i \in F} N^i$; i.e. $x \in \left(\sum_{i \in F} N^i \right)_m$. It follows that

$$R_j x \subseteq \left(\sum_{i \in F} N^i \right)_{m-j} \subseteq \left(\sum_{i \in F} N^i \right)_0$$

(due to $\sum_{i \in F} N^i$ a subfragment of M). Since $\sum_{i \in F} N^i$ is nontrivial, we have that

$$x \in \left(\sum_{i \in F} N^i \right)_i \subseteq K_i.$$

This shows that K is nontrivial. Since

$$d_M(x) = d_{N(F)}(x) \leq d_K(x) \leq d_M(x),$$

whenever $x \in \sum_{i \in F} N^i := N(F)$, and $x \in K_0$, it follows that

$$d_M(x) = d_K(x)$$

for all $x \in K_0$. Therefore K is strict. □

Theorem 12. *Let M be a left R -fragment, and $\{N^i\}_{i \in I}$ be a family of strict subfragments of M . Then*

$$\sum_{i \in I} N^i = \bigcup \left\{ \sum_{i \in F} N^i : F \text{ is a finite subset of } I \right\}.$$

Proof. By Lemma 5,

$$L\left(\bigcup_{i \in I} N_0^i\right) = \bigcup_{i \in I} L(N_0^i) = \bigcup_{i \in I} N_0^i;$$

and hence

$$\prec \bigcup_{i \in I} N_0^i \succ_0 = \left(\sum_{i \in I} N^i \right)_0 = \sum_{i \in I} N_0^i = \bigcup \left\{ \sum_{i \in F} N_0^i : F \text{ is a finite subset of } I \right\}.$$

In Lemma 11, we have seen that $K := \bigcup \left\{ \sum_{i \in F} N^i : F \text{ is a finite subset of } I \right\}$, with the given chain, is a strict subfragment of M . From the definition of the fragment sum of the family $\{N^i\}_{i \in I}$, we deduce that $\sum_{i \in I} N^i := N$

is also a strict subfragment of M . Since $K_0 = N_0$, and $d_N(x) = d_M(x) = d_K(x)$ for all $x \in N_0 (= K_0)$, it follows that $K = N$. \square

Theorem 13. *Let M be a left R -fragment. Then M is finitely spanned if and only if for every family $\{N^i\}_{i \in I}$ of strict subfragments of M ,*

$$\sum_{i \in I} N^i = M \text{ implies } \sum_{i \in F} N^i = M,$$

for some finite subset F of I .

Proof. Let X be a finite spanning set for M , and let $\{N^i\}_{i \in I}$ be a family of strict subfragments of M , with $\sum_{i \in I} N^i = M$. It follows that $X \subseteq \sum_{i \in F} N_0^i$ for some finite subset F of I . Hence $M = \langle X \rangle \subseteq \sum_{i \in F} N^i$, and therefore $M = \sum_{i \in F} N^i$.

It is clear that $M = \sum_{x \in M_0} \langle x \rangle$. Thus, by assumption, $M = \sum_{x \in X} \langle x \rangle = X$ for some finite subset X of M_0 ; i.e. M is finitely spanned. \square

Theorem 14. *Let M be a left R -fragment. If for each i , M_i is finitely generated as an R_0 -module, then every improper subfragment of M is finitely spanned.*

Proof. Without loss of generality we may show that M is finitely spanned. Let $M_0 = \sum_{i=1}^n R_0 x_i$. We claim that M as an R -fragment is spanned by X , where $X = \{x_1, x_2, \dots, x_n\}$. To this end, let $m \in M_k$. Then $m = \sum_{j=1}^s r_j y_j$, where $\{y_1, y_2, \dots, y_s\}$ is a generating set for M_k as an R_0 -module, and $r_j \in R_0$. But for each $i = 1, 2, \dots, s$, we have

$$y_i = \sum_{j=1}^n \beta_{ij} x_j,$$

where $\beta_{ij} \in R_0$. Hence

$$m \in \left\{ \sum_{i=1}^n y_i : y_i \in L(X), d_M\left(\sum_{i=1}^n y_i\right) \geq k \right\};$$

and therefore

$$M_k = \left\{ \sum_{i=1}^n y_i : y_i \in L(X), d_M\left(\sum_{i=1}^n y_i\right) \geq k \right\}; \quad k = 0, 1, \dots$$

Therefore we have our claim. □

4. FACTOR FRAGMENTS

Just as for modules, there is a factor fragment of a left R -fragment with respect to each of strict subfragments. Let M be a left R -fragment and let N be a strict subfragment. Then it is easy to see that the chain of abelian factor groups

$$M_0/N_0 \supseteq (M_1 + N_0)/N_0 \supseteq \dots \supseteq (M_n + N_0)/N_0 \supseteq \dots$$

is a left R -fragment relative to the left multiplication defined via

$$r(x + N_0) = rx + N_0$$

for all $x \in M_n$, $x \in M_n$, and $r \in R_m$, where $m \leq n$.

The resulting fragment (denoted by M/N), is called the left R -factor fragment of M relative to N . Since N is a strict subfragment, then the given left multiplication is well defined. In fact if $f_n : M_n/(N_0 \cap M_n) \rightarrow (M_n + N_0)/N_0$ are the natural abelian group isomorphisms ($n = 0, 1, 2, \dots$), where $f_0 = 1$, then the sequence of monomorphisms

$$M_0/N_0 \xleftarrow{g_1} M_1/N_1 \xleftarrow{g_2} M_2/N_2 \xleftarrow{g_3} \dots M_n/N_n \xleftarrow{g_{n+1}} \dots,$$

where $g_n = (f_{n-1}^{-1} |_{L_n})f_n$ and $L_n := (M_n + N_0)/N_0$, gives rise to the chain

$$M_0/N_0 \supseteq g_1(M_1/N_1) \supseteq g_1g_2(M_2/N_2) \supseteq \dots \supseteq g_1g_2 \dots g_n(M_n/N_n) \supseteq \dots$$

Hence we may consider the chain of the factor R -fragment of M relative to N as:

$$M_0/N_0 \supseteq M_1/N_1 \supseteq M_2/N_2 \supseteq \dots \supseteq M_n/N_n \supseteq \dots,$$

where the left multiplication is give by:

$$r(m + N_n) = rm + N_{n-i},$$

for all $m + N_n \in M_n/N_n$ and $r \in R_i$ where $i \geq n$.

Lemma 15. *Let M be a left R -fragment spanned by a subset X , and let N be a strict subfragment of M . Then M/N is spanned by $\bar{X} = \{x + N_0 : x \in X\}$.*

Proof. From the definition of the left multiplication of R on M/N , we have that

$$r_n(r_{n-1}((r_0x))) + N_0 = r_n(r_{n-1}(\dots(r_0(x + N_0)))),$$

for all $x \in X$, n is a non negative integer, and $r_i \in R_i$; where $i \leq d_M(r_{i-1}(\dots(r_0x)))$, and $i = 0, 1, \dots, n$. Hence, for all $k = 0, 1, \dots$, we have that

$$\begin{aligned} (M/N)_k &= (M_k + N_0)/N_0 = \left[\left\{ \sum_{i=1}^n y_i : y_i \in L(X), d_M\left(\sum_{i=1}^n y_i\right) \geq k \right\} + N_0 \right] / N_0 \\ &= \left\{ \sum_{i=1}^n (y_i + N_0) : y_i \in L(X), d_M\left(\sum_{i=1}^n y_i\right) \geq k \right\} = \prec \bar{X} \succ_k. \end{aligned}$$

□

Corollary 16. *Let M be a finitely spanned left R -fragment, and let N be a strict subfragment of M . Then M/N is finitely spanned.*

Theorem 17. *Let M be a left R -fragment, and let N be a strict subfragment of M . If M/N and N are finitely spanned, then M is finitely spanned.*

Proof. Let $\{\alpha_1, \alpha_2, \dots, \alpha_n\}$ be a spanning set of M/N , and $\{\beta_1, \beta_2, \dots, \beta_m\}$ be a spanning set of N . We claim that M is spanned by

$$X := \{x_1, x_2, \dots, x_n, \beta_1, \beta_2, \dots, \beta_m\},$$

where $x_i \in M_0$ such that $x_i + N_0 = \alpha_i$ ($i = 1, 2, \dots, n$). Now let $m \in M_k$, it follows that $m + N_0 = \sum_{i=1}^s t_i$, $t_i \in L(\{\alpha_1, \alpha_2, \dots, \alpha_n\})$, and $d_{M/N}(\sum_{i=1}^s t_i) \geq k$. Hence $m + N_0 = (\sum_{i=1}^s b_i) + N_0$, where $b_i \in L(\{x_1, x_2, \dots, x_n\})$, and $d_M(\sum_{i=1}^s b_i) \geq k$. Since N is a strict subfragment of M , we have that $d_N(m - \sum_{i=1}^s b_i) \geq k$. Hence $m - \sum_{i=1}^s b_i$, as an element of N_k , can be written as $m - \sum_{i=1}^s b_i = \sum_{i=1}^\lambda c_i$, where $c_i \in L(\{\beta_1, \beta_2, \dots, \beta_m\})$. Therefore

$$m = \sum_{i=1}^s b_i + \sum_{i=1}^\lambda c_i \in \prec \{\{x_1, x_2, \dots, x_n, \beta_1, \beta_2, \dots, \beta_m\}\} \succ_k;$$

i.e.

$$M_k = \left\{ \sum_{i=1}^n y_i : y_i \in L(X), d_M(\sum_{i=1}^n y_i) \geq k \right\}, \quad k = 1, 2, \dots$$

Thus we have our claim. □

5. HOMOMORPHISMS OF FRAGMENTS

Let M and N be two left R -fragments. By a fragment homomorphism $f : M \rightarrow N$ we mean a function $f : M_0 \rightarrow N_0$ such that the following two conditions are satisfied:

1. $d_M(m) \leq d_N(f(m))$, for all $m \in M_0$,
2. $f(rm_1 + sm_2) = rf(m_1) + sf(m_2)$, for all $m_1, m_2 \in M_0$ and $r \in R_i, s \in R_j$, with $i \leq d_M(m_1), j \leq d_M(m_2)$.

Let M and N be two left R -fragments, and let $f : M \rightarrow N$ be a fragment homomorphism. Then the chain

$$f(M_0) \supseteq f(M_1) \supseteq f(M_2) \supseteq \dots \supseteq f(M_n) \supseteq \dots$$

forms a subfragment of N . This subfragment is called the fragment image of f and is denoted by $\text{Im}f$. If K is a subfragment of the fragment N , then the chain

$$f^{-1}(K_0) \supseteq f^{-1}(K_1) \cap M_1 \supseteq f^{-1}(K_2) \cap M_2 \supseteq \dots \supseteq f^{-1}(K_n) \cap M_n \supseteq \dots$$

is a subfragment of M . It is called the fragment inverse image of K under f , and is denoted by $f^{-1}(K)$. The kernel of f (denoted by $\text{Ker}f$) is given by $f^{-1}(0)$, where 0 is the zero subfragment of N . f is called an epimorphism in case of the fragment image $\text{Im}f$ and N , as fragments, are equal. It is called a monomorphism in case of the subfragment $\text{Ker}f$ of M is the zero fragment. A homomorphism which is monomorphism and epimorphism is called an isomorphism.

Proposition 18. *Let M and N be two left R -fragments, and let $f : M \rightarrow N$ be a fragment homomorphism. If K is a strict subfragment of N , then $f^{-1}(K)$ is a strict subfragment of M .*

Proof. It is clear that

$$\begin{aligned} f^{-1}(K_0) \cap M_i &= f^{-1}(K_0) \cap f^{-1}(N_i) \cap M_i \\ &= f^{-1}(K_0 \cap N_i) \cap M_i = f^{-1}(K_i) \cap M_i = f^{-1}(K_i) \end{aligned}$$

(due to K strict), for each $i = 0, 1, 2, \dots$; i.e. $f^{-1}(K)$ is a strict. □

Corollary 19. *Let M and N be two left R -fragments, and let $f : M \rightarrow N$ be a fragment homomorphism. Then $\text{ker}f$ is a strict subfragment.*

Proposition 20. *Let M and N be two left R -fragments, and let $f : M \rightarrow N$ be a fragment homomorphism. If M is spanned by a subset X , then $\text{Im}f$ is spanned by $f(X)$. Moreover if $g : M \rightarrow N$ is also a fragment homomorphism, then $f = g$ if and only if $f(x) = g(x)$ for all $x \in X$.*

Proof. It is clear that $f(r_n(r_{n-1} \dots ((r_0 x)))) = r_n(r_{n-1} \dots ((r_0 f(x))))$, for each $r_n(r_{n-1} \dots ((r_0 x))) \in L(X)$. It follows that

$$\begin{aligned} f(M_k) &= f \left(\left\{ \sum_{i=1}^n y_i : y_i \in L(X), d_M \left(\sum_{i=1}^n y_i \right) \geq k \right\} \right) \\ &= \left\{ \sum_{i=1}^n f(y_i) : y_i \in L(X), d_M \left(\sum_{i=1}^n y_i \right) \geq k \right\} \\ &= \left\{ \sum_{i=1}^n t_i : t_i \in L(f(X)), d_{f(M)} \left(\sum_{i=1}^n t_i \right) \geq k \right\}, \end{aligned}$$

for each $k = 0, 1, 2, \dots$. This shows that $\text{Im} f = \prec f(X) \succ$.

Now let $f(x) = g(x)$ for all $x \in X$. It is easy to check that the chain

$$K_0 \supseteq K_1 \supseteq K_2 \supseteq K_n \supseteq \dots, \quad \text{where } K_i = \{m \in M_i : f(m) = g(m)\},$$

is a strict subfragment of M . Since X is contained in K , we have that $K = M$; and hence $f = g$.

The converse is obvious. □

Corollary 21. *Homomorphic image of a finitely spanned fragment is finitely spanned.*

1. Anderson F. W. and Fuller K. R., *Rings and categories of modules*, Springer Verlag, New York, (1974).
2. Nawal S. Mahmoud, *Fragmented Structures over filtered rings and applications to valuation theory*, Ph. D. Thesis, Faculty of Science, Ain Shams University, Cairo, Egypt, (1994).
3. Nawal S. Mahmoud, and F. Van Oystaeyen, *An introduction to fragmented Structures over filtered rings*, Comm. Alg. **23**(3), (1995), 975–993.

M. A. Kamal, Ain Shams University, Faculty of Education, Mathematics Department, Roxy Cairo, Egypt,
e-mail: mahmoudkamal333@hotmail.com

N. S. Mahmoud, Ain Shams University, Faculty of Education, Mathematics Department, Roxy Cairo, Egypt,
e-mail: nawalsayed2003@yahoo.com